



قائمة الاسئلة

استعاضة صناعية المستوى الرابع - طب وجراحة الفم والاسنان

د. أحمد حويله

- 1) Increased V.D.O can cause the following, except:
 - 1) - Sensation of bulky dentures
 - 2) - Loss of biting power
 - 3) + Poor esthetic like presence of excessive wrinkles
 - 4) - Increased rate of residual ridge resorption
- 2) Regarding the adjusting of bite rims in the frontal plane, which of the following statements is correct?
 - 1) + When there is no sufficient support anteriorly, philtrum becomes flattened and the commissure drops
 - 2) - The anterior border of the occlusion rim should be parallel to the interpupillary line
 - 3) - The anterior segment of bite rim should be inclined anteriorly by 10-15 degrees
 - 4) - The posterior part of the occlusion rim should be parallel to the camper's line
- 3) Regarding the thickness of the alveolar mucosa, which of the following forms an ideal cushion for the basal seat of the denture?
 - 1) + Class I
 - 2) - Class II
 - 3) - Class III
 - 4) - Thick denture-supporting tissues
- 4) Regarding saliva, which of the following can cause gag reflex with denture wearing and tends to displace denture?
 - 1) - Watery saliva
 - 2) - Reduced saliva flow
 - 3) + Viscous saliva
 - 4) - Hyper salivation
- 5) For Bony undercuts, the following statement are true, except:
 - 1) + In the final impression, zinc oxide eugenol material should be used
 - 2) - Prominent bilateral undercuts must be corrected surgically
 - 3) - Relief in the denture over small bilateral or unilateral undercuts can be done
 - 4) - A small unilateral undercut can be solved by changing the denture path of insertion
- 6) "....." offers the best maxillary denture vertical support & retention but can be easily dislodged by a laterally, anteriorly directed forces or rotating forces; so, it considered unfavorable
 - 1) - Class II inter-arch distance
 - 2) - Low round alveolar ridge
 - 3) - Medium tuberosity
 - 4) + Shallow or flat hard palate
- 7) Drugs that may cause xerostomia which adversely affects denture retention, include:
 - 1) - Antihypertensive
 - 2) - Antidepressant
 - 3) - Antipsychotic
 - 4) + All of them
- 8) Regarding the acceptance and tolerance of the patient for prosthetic service, one pair of denture every 1-2 years is considered fair for prognosis
 - 1) + .TRUE
 - 2) - .FALSE
- 9) Gender might be considered a risk factor for masticatory performance of CD wearers
 - 1) + .TRUE





- 2) - .FALSE
- 10) "....." is the assessment of the patient's individual, local, and general conditions by the dentist to prepare the treatment planning
- 1) - Diagnosis
 - 2) - Treatment planning
 - 3) + Examination
 - 4) - All of them
- 11) Which of the following is the main disadvantages of thermoplastic compound impression material?
- 1) + It can't reproduce the undercuts
 - 2) - It can cause irritation for the oral tissues
 - 3) - Difficulties when handled
 - 4) - It is dimensionally unstable
- 12) Impression materials used for making primary impression in complete edentulous patients include the following, except:
- 1) - Alginate
 - 2) + Zinc oxide eugenol
 - 3) - Putty silicon
 - 4) - Modeling compound
- 13) During the border molding of upper arch, moving the patient's cheek outward, downward and inward, then asked the patient to open his mouth widely to record:
- 1) + Posterior extent of the denture
 - 2) - Labial flange
 - 3) - Buccal flange
 - 4) - Shape and thickness of the buccal vestibules
- 14) To assure a correct determination of VDO in fully edentulous people, we can use:
- 1) - Former dentures
 - 2) - Pre-extraction records
 - 3) + Using several methods to reach a precise VDO
 - 4) - Phonetics & esthetics tests
- 15) Regarding Willis method of recording V.D.R, all the following are true, except:
- 1) - At rest these measurements must be equal
 - 2) - The distance is measured using a special device called willi's gauge
 - 3) - It is designed to measure the distance from the canthus of the eye to the commissure of the mouth & the distance from the lower border of the nose to the gnathion
 - 4) + It can be used in asymmetric face
- 16) The accuracy of maxillo-mandibular relation record is affected by the following, except:
- 1) + Treatment outcome prognosis
 - 2) - Retention of the record bases
 - 3) - Stability of record base
 - 4) - Rigidity of record base
- 17) Regarding soft palate slopes, which one of the following forms a good posterior seal for denture?
- 1) + Class I
 - 2) - Class II
 - 3) - Class III
 - 4) - Class IV
- 18) Regarding the phonetics test, asking the patient to repeat the name "emma" will help to record and assess:
- 1) + V.D.R
 - 2) - V.D.O
 - 3) - Centric relation





- 4) - Smile line
- 19) All of the following statements, regarding the primary impressions, are true, except:
- 1) + It is made with alginate or compound in plain stock tray
 - 2) - It is made for the purpose of diagnosis, treatment planning, and for the construction of individual tray
 - 3) - The method for making it should be based on the basic principles of maximum area coverage and intimate contact
 - 4) - It is made by using stock tray and cheap material
- 20) According your clinical experience, which of the following considers the best treatment option for mandibular complete edentulous patients?
- 1) - Conventional complete denture
 - 2) + Implant retained over denture
 - 3) - Implant retained fixed denture
 - 4) - All of the above

