



قائمة الاسئلة

اطقم جزئيه 6 -الرابع -فني الاسنان- كلية طب وجراحة الفم والاسنان- درجة الامتحان (20)

د / نهله الجحدري

- 1) We can use Flexible dentures in cases of conventional acrylic sensitivity
  - 1) ☒ TRUE.
  - 2) ☐ FALSE.
- 2) ..... resist occlusal wear and are well suited for maintaining vertical dimension during provisional restorative therapy.
  - 1) ☒ Acetal resins
  - 2) ☐ Thermoplastic polyester
  - 3) ☐ Thermoplastic polyamide (nylon)
  - 4) ☐ ALL
- 3) Advantages of flexible denture include
  - 1) ☐ the denture base adapts well in the undercut areas
  - 2) ☐ Aesthetic translucency
  - 3) ☐ strong that it can be made very thin which make it comfortable to wear and esthetically pleasing
  - 4) ☒ All
- 4) while setting the teeth in the flexible denture we should consider..... To prevent debonding of the teeth from the denture base
  - 1) ☐ Sufficient height of the selected teeth
  - 2) ☐ Mechanical undercuts (diatorics)
  - 3) ☒ All
  - 4) ☐ none
- 5) The flexible denture base material has a unique property that it.....chemically bonds with any of the acrylic resin or porcelain
  - 1) ☐ does
  - 2) ☒ does not
- 6) Extreme caution is necessary when processing the flexible dentures as
  - 1) ☐ proper ventilation,masks, and vaccum systems
  - 2) ☒ all
  - 3) ☐ none
- 7) No repair or relining is feasible with flexible dentures
  - 1) ☒ TRUE.
  - 2) ☐ FALSE.
- 8) Contraindications of flexible denture include
  - 1) ☐ Patients with insufficient inter arch space
  - 2) ☐ Flat flabby ridges with poor soft tissue support
  - 3) ☐ Bilateral free-end distal extension on maxilla with extremely atrophied alveolar ridges
  - 4) ☒ all
- 9) After RPD insertion,tissue blanching and inflammation or ulceration seen when the beading on the cast done
  - 1) ☒ too deep
  - 2) ☐ too shallow
  - 3) ☐ all
  - 4) ☐ none
- 10) Causes of incomplete casting of the metal framework are
  - 1) ☐ metal too cold when cast
  - 2) ☐ improper spruing
  - 3) ☐ gas trapped in mold



- 4) ☒ All
- 11) If too much grinding is required to correct occlusion in RPD insertion, the solution is
- 1) - spot grinding using the articulating paper
  - 2) - the use of articulator in adjusting occlusion is preferred
  - 3) ☒ remove the denture base and the artificial tooth , and reconstruct a new denture base
  - 4) - NONE
- 12) Causes of open bite after the insertion of RPD
- 1) - when the investment material is still soft and the acrylic is pressed in the flask
  - 2) - too great or too rapid pressure applied during initial packing might force the teeth into the plaster
  - 3) - failure to close the flask completely at final closure
  - 4) ☒ ALL
- 13) Causes of no retention of the RPD
- 1) ☒ improper placement of the retentive clasp in the frame work of the partial denture
  - 2) - improper placement of the rest in the frame work of the partial denture
  - 3) - improper placement of the major connector in the frame work of the partial denture
  - 4) - improper placement of the reciprocal clasp in the frame work of the partial denture
- 14) A digital fabrication method(CAD-CAM) has various advantages as
- 1) - electronic surveying
  - 2) - determine the path of insertion and removal and control the undercut amount through computation
  - 3) - the design data may be saved for future
  - 4) ☒ All
- 15) Pathways of manufacturing of a denture framework are
- 1) - manufacturing of metal framework directly by selective laser melting technology
  - 2) - fabricate a sacrificial pattern made of resin
  - 3) - designing and wax pattern fabrication then investing and casting procedures
  - 4) ☒ ALL
- 16) This involve physically removing or subtracting materials with cutting tools to achieve the selected shape
- 1) ☒ Subtractive method
  - 2) - Additive method
  - 3) - ALL
  - 4) - NONE
- 17) In maxillofacial partial dentures
- 1) - Multiple rests can be used to improve the stability
  - 2) - Maximum retention, support and stability can be obtained by utilizing the defect
  - 3) - Ideal contours for retention, guiding planes and occlusal rests by full crowns of the abutments teeth
  - 4) ☒ ALL
- 18) In maxillofacial partial denture,the framework and the moulded resin is used as special tray for making the final impression
- 1) ☒ TRUE.
  - 2) - FALSE.
- 19) In maxillofacial RPD,remaining teeth on the defective side shifts the fulcrum posteriorly the indirect retainers should be placed ..... as possible
- 1) ☒ anteriorly
  - 2) - posteriorly
- 20) In cases of speech aid prostheses the major connector functionally extended palatally to restore deficient areas of soft palate
- 1) ☒ TRUE.
  - 2) - FALSE.