



قائمة الاسئلة 2025-04-30 04:12

زراعة اسنان2-الرابع-تكنولوجيا صناعة الاسنان-كلية طب وجراحة الفم والاسنان-درجة الامتحان (20)

د. محمد عبدالعزيز البيلي

- 1) With fixed partial dentures, it has been recommended that ,,,, of mesio-distal space be available when placing 2 implants.
 - 1) - 21 mm
 - 2) - 23 mm
 - 3) - 24 mm
 - 4) ☒ None
- 2) The surgical anatomy of the maxilla and mandible have that may cause injury to blood vessels and nerves.
 - 1) ☒ 4 functional zones
 - 2) - 5 functional zones
 - 3) - 6 functional zones
 - 4) - 7 functional zones
- 3) What are the benefits of Impression in implant dentistry.
 - 1) - Position.
 - 2) - Depth.
 - 3) - Axis/Angulation
 - 4) ☒ All
- 4) In some cases, the molar must replace by two implants and not only one wide implant to
 - 1) - Increase the torque on the part not supported by the fixture.
 - 2) ☒ Decrease the torque on the part not supported by the fixture.
 - 3) - All
 - 4) - None
- 5) Adequate vertical space must be available into which a crown can be placed for the best esthetic & structural forms.
 - 1) ☒ Inter-arch Space.
 - 2) - Intra arch distance
 - 3) - All
 - 4) - None
- 6) In esthetically demanding situations the implant is placed below the gingival crest
 - 1) - 1 - 2 mm
 - 2) ☒ 2 - 3 mm
 - 3) - 3 - 4 mm
 - 4) - None
- 7) Use the with open tray impression technique.
 - 1) ☒ Bick- up.
 - 2) - Transfer
 - 3) - Gingival former
 - 4) - All
- 8) Full arch restorations should not be considered on less than implants in the maxilla and in the mandible.
 - 1) - Five - Six
 - 2) - Five - Seven
 - 3) ☒ Seven - Five
 - 4) - Seven -Six
- 9) records the density, thickness of bone & bony contours.



- 1) - Panoramic radiography
 - 2) - CT scan
 - 3) ☒ CBCT
 - 4) - None
- 10) Using healing cap or interim abutment to
- 1) - Maintain proper emergence profile.
 - 2) - Esthetics and healthy peri-implant tissue.
 - 3) ☒ All
 - 4) - None
- 11) is a replica of the entire dental implant not intended for the human implantation
- 1) ☒ Implant analogue.
 - 2) - Impression coping
 - 3) - Gingival former
 - 4) - None
- 12) Use the with closed tray impression technique.
- 1) - Bick- up.
 - 2) ☒ Transfer
 - 3) - Gingival former
 - 4) - All
- 13) Used when there is limited inter arch space and tendency to gaging.
- 1) - Open tray impression technique
 - 2) ☒ Closed tray impression technique.
 - 3) - All
 - 4) - None
- 14) Principles of implant-tooth connection: may be.
- 1) - Non rigid connection.
 - 2) - Multiple teeth (abutments) incorporated within the long span implant restoration.
 - 3) - Telescopic crown should be used to cover the natural tooth to prevent decay if loosening occurs.
 - 4) ☒ All
- 15) Sharp caliper is pushed through the tissues till bone contact, to determine at the proposed implant sites.
- 1) ☒ The soft tissue thicknesses.
 - 2) - The hard tissue thicknesses.
 - 3) - All
 - 4) - None
- 16) Surgical stents are made of clear resin preferably to.
- 1) - Locate the implant within restoration contour.
 - 2) - Align implant with long axis of completed restoration.
 - 3) - Identify the level of CEJ or tooth emergence from soft tissue
 - 4) ☒ All
- 17) If implant placement too far buccally will lead to
- 1) ☒ Breakdown of buccal plate of bone.
 - 2) - Breakdown of lingual plate of bone.
 - 3) - All
 - 4) - None
- 18) Implant should be away from adjacent natural teeth.
- 1) - 1 mm
 - 2) ☒ 1.5mm
 - 3) - 2 mm



- 4) - None
- 19) Implant should be away from adjacent implant.
- 1) - 1 mm
- 2) - 1.5mm
- 3) + 3 mm
- 4) - None
- 20) Mounted casts are used to.
- 1) - Study the remaining dentition.
- 2) - Evaluate the residual bone.
- 3) - Analyse maxillo-mandibular relationships
- 4) + All