



قائمة الاسئلة

امتحان نهاية الفصل الدراسي الأول - للعام الجامعي 1446 هـ - كلية العلوم :: ميكروبيولوجيا تشخيصية متقدمة - (434102) - المستوى الرابع
أنس أحمد حسين المحبشي

- Meningitis may occur due to infection by (1)
- staph aureus - (1)
 - streptococcus pneumonia + (2)
 - Bacillus anthracis - (3)
 - None - (4)
- Thayer martin agar is a selective media for (2)
- S.aureus - (1)
 - B anthracis - (2)
 - N.meningitidis + (3)
 - None - (4)
- Manitol salt Agar is a selective media for (3)
- S.pyogenes - (1)
 - M.tuberculosis - (2)
 - N.meningitidis - (3)
 - S.aureus + (4)
- All are from Bacteria that may cause Food poisoning Except (4)
- C.botulinum - (1)
 - Staph.aureus - (2)
 - Strept.pyogens + (3)
 - B.cerus - (4)
- Gas gangrene is mainly occure due to infection by (5)
- M.tuberculosis - (1)
 - Staph.aureus - (2)
 - Bacillus anthracis - (3)
 - C.perfringens + (4)
- In the Lab, we need calibrated loop in case of (6)
- Stool culture - (1)
 - Pus culture - (2)
 - C.S.F culture - (3)
 - Urine culture + (4)
- The time required for Blood culture in routin media is (7)
- 4-6 weeks + (1)
 - 1-2 weeks - (2)
 - 1-2 months - (3)
 - None - (4)
- Undulent (maltic) fever is may occur due to infection (8)
- Salmonella.typhi - (1)
 - Escherishia coli - (2)
 - Pseudomonas aeroginusa - (3)
 - None + (4)
- In case of Bacterial meningitis the rate of Glucose in the C.S.F sample is (9)
- decreased + (1)
 - Increased - (2)
 - Normal - (3)
 - None - (4)
- The sample used for diagnosis of Brucella in the Lab is (10)





C.S.F culture	-	(1)
Urine culture	-	(2)
Pus culture	-	(3)
Blood culture	+	(4)
Ogawa media and L-J media is a selective media for		(11)
S.pyogenes	-	(1)
M.tuberculosis	+	(2)
N.menigitidis	-	(3)
S.aureus	-	(4)
Most Skin Infectionis occur due to Infection by		(12)
Strapt.pyogens	-	(1)
staph aureus	+	(2)
C.perfringens	-	(3)
Listeria monocytogens	-	(4)
Rheumatic heart disease is mainly occue due to infection by		(13)
Strapt.pyogens	+	(1)
staph aureus	-	(2)
C.perfringens	-	(3)
Coryni bacterium	-	(4)
Indicate for Bacterial infection in the Urine		(14)
Protien +ve	-	(1)
Glucose +ve	-	(2)
Blood+ve	-	(3)
Nitrite +ve	+	(4)
Diphesic Media (Agar/broth) is used in		(15)
C.S.F culture	-	(1)
Urine culture	-	(2)
Pus culture	-	(3)
Blood culture	+	(4)
Bacillary dysentry is may occure due to infection by		(16)
Salmonella	-	(1)
Shigella	+	(2)
cholera	-	(3)
Clostridium	-	(4)
Infection by Clost.perfringens is more sever in patients		(17)
Young children	-	(1)
Diabetic	+	(2)
old age	-	(3)
meningitis	-	(4)
The best media for Stool culture is		(18)
XLD	+	(1)
Macconcky	-	(2)
chocolate	-	(3)
CLED	-	(4)
The best media for Urine culture is		(19)
XLD	-	(1)
Macconcky	-	(2)
chocolate	-	(3)
CLED	+	(4)





- Ziehl-Neelsen smear should be done when (20)
- Neisseria suspected - (1)
 - Cholera suspected - (2)
 - Tuberculus Bacilli suspected + (3)
 - All - (4)
- all these bacteria causing Bacteremia Except (21)
- Staph.aureus - (1)
 - Srept.pyogens - (2)
 - Haemophilus influenzae - (3)
 - Corynebacterium diphtheriae + (4)
- All these bacteria causing Urinary tract infection Except (22)
- Staph.saprophyticus - (1)
 - Srept.pyogens + (2)
 - Escherishia coli - (3)
 - Pseudomonas aeruginosa - (4)
- The media used for cultring of Fungi in the Lab is (23)
- XLD - (1)
 - CLED - (2)
 - Sabaroud Dextrose agar + (3)
 - TCBC - (4)
- The best media for culturing Pus sample is (24)
- Blood agar and MacConkey + (1)
 - Chocolate Agar - (2)
 - S-S Agar - (3)
 - None - (4)
- The most bacteria that causing Throat Infection is (25)
- Strept. pyogenes + (1)
 - Staph.aureus - (2)
 - Escherishia coli - (3)
 - All - (4)

