



قائمة الاسئلة 2025-04-09 06:25

علم الديدان الطفيلية - (414202) - المستوى الرابع - تخصص علم الاحياء - كلية العلوم - الفترة الثالثة - درجة الامتحان (50)

ابتهاج حسن العجمي

- 1) Infective stage of *S. hematobium* is
 - 1) - Miracidium
 - 2) + Cercaria
 - 3) - Worm
 - 4) - Eggs
- 2) All these worms are endoparasitic
 - 1) - Turbellaria
 - 2) - insects
 - 3) + Cestodae
 - 4) - Monogenoidea
- 3) The mouth and alimentary canal are absent in
 - 1) + Dipylidium caninum
 - 2) - S. japonicum
 - 3) - Clonorchis
 - 4) - Heterophyes
- 4) The egg shape is small with lateral spine in
 - 1) - S. intercalatum
 - 2) - S. mansoni
 - 3) - S. haematobium
 - 4) + S. japonicum
- 5) Oviparous, it means
 - 1) - producing larvae
 - 2) - laying eggs containing larvae
 - 3) - laying larvae
 - 4) + laying eggs
- 6) Mode of infection is Penetration of skin as
 - 1) + Ancylostoma
 - 2) - Enterobius
 - 3) - Dracunculus
 - 4) - Ascaris
- 7) Tissues nematode parasite as
 - 1) - Trichuris trichiura
 - 2) + Larva migrans
 - 3) - Strongyloides
 - 4) - Hook worm
- 8) The mouth with three lips in
 - 1) - whip-worm
 - 2) - pin worm
 - 3) - Beef tape worm
 - 4) + Giant intestinal nematode
- 9) Biomphalaria is intermediate host snail of
 - 1) - S. intercalatum
 - 2) + S. mansoni
 - 3) - S. haematobium
 - 4) - S. japonicum





- 10) Dog tape worm infection is
- 1) - *Taenia saginata*
 - 2) - *Diphyllobothrium latum*
 - 3) + *Dipylidium caninum*
 - 4) - *Echinococcus granulosus*
- 11) The anterior, undifferentiated segmented
- 1) + Immature segments
 - 2) - Mature segments
 - 3) - Gravid proglottis
 - 4) - Genital pores
- 12) Hydatid cyst is infective stage for
- 1) + man
 - 2) - dog
 - 3) - rat
 - 4) - insect
- 13) Dwarf tape worm is
- 1) + *Hymenolepis nana*
 - 2) - *Hymenolepis diminuta*
 - 3) - *D. latum*
 - 4) - *T. Sagenata*
- 14) They have two intermediates
- 1) - *Strongyloides stercoralis*
 - 2) + *Dicrocoelium dendriticum*
 - 3) - *Ascaris*
 - 4) - *S. mansoni*
- 15) The esophagus takes different shapes in different species in
- 1) + Nematoda
 - 2) - Cestodae
 - 3) - Monogeneoidea
 - 4) - Trematoda
- 16) Hydatid cysts are found in liver of man
- 1) + 70%,
 - 2) - 20%
 - 3) - 10%
 - 4) - 1%
- 17) whip-worm is
- 1) - *Ancylostoma*
 - 2) - *Enterobius vermicularis*
 - 3) - *Ascaris lumbricoides*
 - 4) + *Trichuris trichiura*
- 18) Gravid segments has a uterus with 7-13 lateral branches in
- 1) - dwarf tape worm
 - 2) + pork tape worm
 - 3) - rat tape worm
 - 4) - broad tape worm
- 19) Worms are tape or ribbon like
- 1) - Turbellaria
 - 2) - Nematoda
 - 3) + Cestodae



4) - Monogenoidea


20)



This segment is

- 1) + Mature
- 2) - Graid
- 3) - Immature
- 4) - Neck

21)



- 1) - Miracidium
- 2) + Cercaria
- 3) - Redia
- 4) - snail

22)



This parasite is

- 1) - Diphyllbothrium latum
- 2) - Dipylidium caninum
- 3) - Hymenolepis diminuta
- 4) + Echinococcus granulosus

23)




This shape called

٢٢

- 1) - Cysticercus cellulosae
- 2) - Coenurus cerebralis
- 3) + Hydatid cyst
- 4) - Pseudophyllidea


24)



[This part refer to

- 1) - Male
- 2) + Female
- 3) - both
- 4) - Larva

25)



This life cycle for

- 1) + Hymenolepis
- 2) - Taenia
- 3) - Dipylidium



4) - Diphyllbothrium