



قائمة الاسئلة 2025-04-20 05:34

علم الطفيليات-(432205)-الثاني-احياء دقيقة-كلية العلوم-درجة الامتحان(50)

أنس أحمد المحبشي

- 1) Presence of mucus with blood in the stool may indicated for infection by
 - 1) - C. parvum
 - 2) - G. lambelia
 - 3) - Toxoplasma
 - 4) + E.histolytica
- 2) The routine sample that be used for diagnosis of Toxoplasosis is
 - 1) + Blood
 - 2) - Urine
 - 3) - Stool
 - 4) - None
- 3) The diagnostic stage in coetaneous Leishmaniasis is
 - 1) - gametocytes
 - 2) - Promastigote
 - 3) - Oocyst
 - 4) + None
- 4) Amoeba present as a normal flora in the intestine is
 - 1) - Entamoeba histolytica
 - 2) - Naegleria fowleri
 - 3) + Entamoeba coli
 - 4) - none
- 5) The diagnostic stage of T.vaginalis is
 - 1) - Cyst
 - 2) - Oocyst
 - 3) + trophozoit
 - 4) - None
- 6) All are correct characters of Leishmania Except
 - 1) - infect macrophage
 - 2) + Non motile
 - 3) - Indirect life cycle
 - 4) - the vector is female sand fly
- 7) Rectum Eczyma may occur due to infection by
 - 1) - Leishmania tropica
 - 2) - Schistosoma mansoni
 - 3) - Ascaris lumbricoides
 - 4) + none
- 8) Appearance of Blood in stool may occur due to infection by
 - 1) - Schistosoma haematobium
 - 2) + Schistosoma mansoni
 - 3) - Taenia saginata
 - 4) - Echinoccuc granolusus
- 9) The drug of choice against Schistosomiasis is
 - 1) - albindazole
 - 2) + praziquantel
 - 3) - Spiramycin
 - 4) - none



- 10) Infection by *Echinococcus granulosus* occur through
- 1) - contaminated cow meat
 - 2) - contaminated Pig meat
 - 3) - contaminated cat feces
 - 4) ☒ contaminated dog feces
- 11) The Incubation period of *Enterobius* egg in the environment is
- 1) - 1-2 week
 - 2) - 2-3 weeks
 - 3) - 1- 2 months
 - 4) ☒ none
- 12) Hydated cyst disease is occur due to infection by
- 1) - *Taenia solium*
 - 2) - *T.saginata*
 - 3) - *S.haematobium*
 - 4) ☒ *Echinococcus granulosus*
- 13) Albendazole treatment for 4 months should be given in case of infection by
- 1) - *H.nana*
 - 2) - cysticercosis
 - 3) ☒ Hydated cyst
 - 4) - *Schistosoma*
- 14) The sample used for diagnosis of infection by *E.granulosus* is
- 1) - C.S.F
 - 2) - Urine
 - 3) - Stool
 - 4) ☒ None
- 15) Cysticercosis in human is occur due to
- 1) ☒ *Taenia solium*
 - 2) - *T.saginata*
 - 3) - *S.haematobium*
 - 4) - *Echinococcus granulosus*
- 16) Pregnant women should be tested for all this tests except
- 1) - Toxoplasma
 - 2) - Rubella
 - 3) - Herpes
 - 4) ☒ *H.nana*
- 17) taenia worm contain from the following
- 1) - Head
 - 2) - Neck
 - 3) - Proglotides
 - 4) ☒ All
- 18) In *T.saginata* Life cycle human is considere
- 1) - intermidated host
 - 2) ☒ definitive Host
 - 3) - All
 - 4) - None
- 19) Delay in children growth may occur in case of infection by
- 1) ☒ *T.saginata*
 - 2) - *H.nana*
 - 3) - *schistosoma*



- 4) - All
- 20) Osphegos bleeding (Varicose) may occur due to infection by
- 1) - T.saginata
 - 2) - H.nana
 - 3) + schistosoma
 - 4) - Echinococcus granolusus
- 21) the most severe species of Plasmodium is
- 1) - P.ovale
 - 2) - P.malariae
 - 3) - P.vivax
 - 4) + None
- 22) The routine sample that be used for diagnosis of Malaria is
- 1) - Urine
 - 2) + Blood
 - 3) - Sputum
 - 4) - None
- 23) Parasite that may causing Abdominal Blockage
- 1) - Schistosoma
 - 2) - Enterobius
 - 3) - Trichuris
 - 4) + Ascaris
- 24) In the life cycle of Echinococcus; human is considered
- 1) - Defenitive host
 - 2) + intermidated host
 - 3) - All
 - 4) - None
- 25) Blader carcinoma may occur due to infection by
- 1) - Taenia solium
 - 2) - T.saginata
 - 3) + S.haematobium
 - 4) - E. granolusus

