



قائمة الاسئلة 2025-05-24 05:45

اسعافات اوليه - كلية الصيدلة - المستوى الثالث-درجة الامتحان ( 50 )

أ.د. عبدالسلام دلاق

- 1) Venous bleeding can be pervented easily.
  - 1) ☒ TRUE.
  - 2) ☐ FALSE.
- 2) External bleeding is very serious than internal bleeding.
  - 1) ☐ TRUE.
  - 2) ☒ FALSE.
- 3) Arterial bleeding is usually dark red color.
  - 1) ☐ TRUE.
  - 2) ☒ FALSE.
- 4) The ingested poisons are the poisons that are breath in
  - 1) ☐ TRUE.
  - 2) ☒ FALSE.
- 5) The second degree burns if infected require skin grafting to heal
  - 1) ☒ TRUE.
  - 2) ☐ FALSE.
- 6) The lower limbs bleeding can be prevented by pressure point on femoral artery.
  - 1) ☒ TRUE.
  - 2) ☐ FALSE.
- 7) Bleeding from the face can be prevented by pressure point on ulnar artery.
  - 1) ☐ TRUE.
  - 2) ☒ FALSE.
- 8) The respiratory rate in adult at bed rest is ranged between 20 - 35/ minute
  - 1) ☐ TRUE.
  - 2) ☒ FALSE.
- 9) Sizures occur as a result of brain tumor.
  - 1) ☒ TRUE.
  - 2) ☐ FALSE.
- 10) Anaphylactic shock may occurs as a result of bacteria.
  - 1) ☐ TRUE.
  - 2) ☒ FALSE.
- 11) Injected poisons are the poisons that are inserted through the skin
  - 1) ☒ TRUE.
  - 2) ☐ FALSE.
- 12) Cardiogenic shock occurs as a result of myocardial infarction.
  - 1) ☒ TRUE.
  - 2) ☐ FALSE.
- 13) Respiration is the process of inspiration and expiration.
  - 1) ☒ TRUE.
  - 2) ☐ FALSE.
- 14) Obstruction by foreign objects or materials is most commonly seen with adult concsiou patient.
  - 1) ☐ TRUE.
  - 2) ☒ FALSE.
- 15) The chest thrusts is a method used to remove foreign objects from an airway with pregnant woman in the late stage of pregnancy.
  - 1) ☒ TRUE.



- 2) - FALSE.
- 16) Bleeding from arm & shoulder can be prevented by pressure point on subclavian artery.
- 1) ☒ TRUE.
- 2) - FALSE.
- 17) Non-Insulin dependent diabetes mellitus usually begins in childhood age
- 1) - TRUE.
- 2) ☒ FALSE.
- 18) Abuse or withdrawal of drugs or alcohol can cause seizures
- 1) ☒ TRUE.
- 2) - FALSE.
- 19) Hypoglycemia occurs as a result of the inability of pancreas to produce enough insulin.
- 1) - TRUE.
- 2) ☒ FALSE.
- 20) Direct force or pressure can fracture a bone at the point of contact.
- 1) ☒ TRUE.
- 2) - FALSE.
- 21) Hypovolemic shock occurs as a result of the following:
- 1) ☒ Loss of blood
- 2) - Myocardial infarction
- 3) - Allergic reaction
- 4) - Bacteria
- 22) Cardiogenic shock occurs as a result of the following :
- 1) - Bacteria
- 2) - Allergic reaction
- 3) - Loss of blood
- 4) ☒ Myocardial infarction
- 23) The second degree burns can be characterized by the following except one:
- 1) - Painful
- 2) - Blisters
- 3) ☒ Generally require 1-3 days to heal without appropriate care
- 4) - Swollen
- 24) The causes of bleeding include the following except one
- 1) - Rptured organs
- 2) - Crushing injuries
- 3) - Ulcer bleeding
- 4) ☒ Constipation
- 25) The bleeding from the head can be prevented by pressure point on
- 1) - Femoral artery
- 2) - Facial artery
- 3) ☒ Temporal artery
- 4) - Radial artery
- 26) The bleeding from the face can be prevented by pressure point on
- 1) - Femoral artery
- 2) ☒ Facial artery
- 3) - Subclavian artery
- 4) - Ulner artery
- 27) The signs and symptoms of Fractures include :
- 1) - Tenderness and pain
- 2) - Deformity of a limb





- 3) - Loss of distal pulse  
4) + All of above
- 28) The reasons for splinting include the following except one:  
1) + To move the injured extremity  
2) - To prevent or reduce the pain  
3) - To prevent damage to soft tissues  
4) - To prevent bleeding
- 29) Patient care with snakebites :include the following except one  
1) - Keep the patient calm  
2) + Move the bitten extremities  
3) - Locate the fang marks and clean this site with soap and water  
4) - Apply a light constricting band above and below the wound
- 30) Patient care with complete airway obstruction: include the following :  
1) - Open the airway  
2) - Provide artificial ventilations for unconscious, nonbreathing patient.  
3) - Remove any foreign object  
4) + All of above
- 31) The signs of internal bleeding include the following except one  
1) - Wound that have penetrated the skull  
2) + Hypertension  
3) - Vomiting or coughing up blood  
4) - Abdominal tenderness & rigidity
- 32) Patient care for absorbed poisons include the following except one::  
1) - Move the patient from the source of the poison or source from patient  
2) + Dry chemicals should be wash directly by soap and water  
3) - Remove all contaminated clothing  
4) - Be on the alert for anaphylactic shock
- 33) The patient care with ingested poisons include the following except one:  
1) + Give anything by mouth if the patient is having a seizure  
2) - Maintain an open airway and transport immediately to the center or hospital.  
3) - For conscious patient's, dilute the poison by having the patient drink are two glasses of water or milk  
4) - Position the conscious patient in semi recumbent position
- 34) The inhaled poisons include the following except one  
1) - Car exhaust  
2) - Burning stoves & furnaces  
3) + Improper prepared food  
4) - Insect sprays
- 35) The first degree burns can be characterized by the following except one  
1) - Painful  
2) - Red in color  
3) - Swollen  
4) + Heal within 1-2 week without of appropriate care
- 36) Septic shock occurs as a result of the following  
1) - Loss of blood  
2) - Myocardial infarction  
3) - Allergic reaction  
4) + Bacteria
- 37) Anaphylactic shock occurs as a result of the following :





- 1) - Bacteria
  - 2) + Allergic reaction
  - 3) - Loss of blood
  - 4) - Myocardial infarction
- 38) The signs & symptoms of insects bites include except one:
- 1) - Noticeable stings or bites on the skin
  - 2) - Localized pain or itching
  - 3) - Numbness in a limb or body part
  - 4) + Normal breathing & pulse
- 39) Hyperglycemia occurs as a result of the following except one:
- 1) - Patient has not taken enough insulin
  - 2) + Patient takes too much insulin
  - 3) - Patient has forgotten to take his insulin
  - 4) - Patient has overeaten
- 40) The causes of siezures include the following:
- 1) - Brain tumor
  - 2) - Head injuries
  - 3) - High fever in children
  - 4) + All of above