



قائمة الاسئلة

التدخل الجراحي والطبي الإكلينيكي - المستوى الثالث - قسم علاج طبيعي - كلية الطب والعلوم الصحية - برامج العلوم الطبية التطبيقية - الفترة د/ وليد أمين عبدالله المخلافي

- 1) One of the following shock is due to low peripheral arteriolar resistance (vasodilatation)
 - 1) Hypovolemic shock
 - 2) Cardiogenic shock
 - 3) Obstructive shock
 - 4) + Neurogenic shock
- 2) To calcutate the extent of burn surface we use
 - 1) Rule of 7
 - 2) Rule of 8
 - 3) + Rule of 9
 - 4) Rule of 10
- 3) The gallbladder is found in the
 - 1) Left hypochondrium
 - 2) + Right hypochondrium
 - 3) Left iliac fossa
 - 4) Right iliac fossa
- 4) Routine preoperative preparation include the following EXCEPT
 - 1) History
 - 2) Examination
 - 3) Informed consent
 - 4) + Brain CT scan
- 5) Presence of blood in the urine is called
 - 1) Pyuria
 - 2) Necroturia
 - 3) + Hematuria
 - 4) Pneumaturia
- 6) All the following are signs and symptoms of compartment syndrome EXCEPT
 - 1) Pulselessness
 - 2) Pain disproportionate to injury
 - 3) Pallor
 - 4) + Intact senation
- 7) All the following are mentioned in the past history EXCEPT
 - 1) Past medical history
 - 2) History of blood transfusion
 - 3) Past surgical history
 - 4) + Chief complain
- 8) The most common injury of the esophagus is
 - 1) Penetrating injury
 - 2) Blunt injury
 - 3) + Iatrogenic injury
 - 4) Malignancy
- 9) The most common cause of acute pancreatitis is
 - 1) Malignancy
 - 2) + GB stones
 - 3) Trauma
 - 4) Drugs



- 10) The lingula (of the lung) is part of the
 - 1) Right upper lobe of the lung
 - 2) Right lower lobe of the lung
 - 3) + Left upper lobe of the lung
 - 4) Left lower lobe of the lung
- 11) The primary cause of death in burnt patients is
 - 1) Constricting eschars
 - 2) + Infection
 - 3) Scar formation
 - 4) Contractures
- 12) Beck's triad of the cardiac tamponade include the following EXCEPT
 - 1) Hypotension
 - 2) Muffled heart sounds
 - 3) Distended neck veins
 - 4) + Decrease air entry
- 13) Empyema means
 - 1) Presence of blood in the pleural cavity
 - 2) + Presence of pus in the pleural cavity
 - 3) Presence of chyle in the pleural cavity
- 14) One of the following statements is FALSE
 - 1) + In acute hepatitis, elective surgery should not be postponed
 - 2) In acute hepatitis, emergency surgery should not be postponed
 - 3) In chronic hepatitis, elective surgery should not be postponed
 - 4) In chronic hepatitis, emergency surgery should not be postponed
- 15) The first stage of wound healing is
 - 1) Remodeling phase
 - 2) + Inflammatory phase
 - 3) Maturation phase
 - 4) Proliferation phase
- 16) The most common site of bilharziasis is the
 - 1) Kidneys
 - 2) Ureters
 - 3) + Urinary bladder
 - 4) Urethra
- 17) Acute cholecystitis means
 - 1) Acute inflammation of the common bile ducts
 - 2) + Acute inflammation of the gallbladder
 - 3) Acute inflammation of the liver
 - 4) Acute inflammation of the spleen
- 18) In humeral shaft fracture there is injury of
 - 1) + Radial nerve
 - 2) Median nerve
 - 3) Ulnar nerve
 - 4) Axillary nerve
- 19) Blood loss in hemorrhagic shock between 30-40% of blood is
 - 1) Class I Hemorrhagic shock
 - 2) Class II Hemorrhagic shock
 - 3) + Class III Hemorrhagic shock
 - 4) Class IV Hemorrhagic shock

4 / 2 الصفحة



- 20) One of the following statements regarding peptic ulcers is TRUE
 - 1) + Duodenal ulcers are more common than gastric ulcers
 - 2) Duodenal ulcers carry more risk for malignancy than gastric ulcers
 - 3) Duodenal ulcers occur in older ages than gastric ulcers
 - 4) Duodenal ulcers occur in the antrum of the stomach
- 21) All the following are true regarding Acute Epidural Hematoma EXCEPT
 - 1) There is usually mild trauma to the brain
 - 2) There is a lucid period
 - 3) Early surgey gives good results
 - 4) + It is due to accumulation of blood between dura and arachnoid matters
- 22) All the following are functions of the skin EXCEPT
 - 1) Protection
 - 2) + Synthesis of vitamin K
 - 3) Synthesis of vitamin D
 - 4) Thermoregulation
- 23) All the following are characteristic of 2nd degree burns EXCEPT
 - 1) They are moist and erythema
 - 2) There are blisters
 - 3) + They are painless
 - 4) They often heal within 3 weeks
- 24) All the following regarding BPH are true EXCEPT
 - 1) + BPH means benign prostatic hypertrophy
 - 2) BPH means benign prostatic hyperplasia
 - 3) BPH is a normal aging process
 - 4) The origin of BPH is the transitional zone
- 25) The ulcers that occur in bed-ridden patients are called
 - 1) + Pressure ulcers
 - 2) Diabetic ulcers
 - 3) Venous ulcers
 - 4) Malignant ulcers
- 26) The most commn type of shoulder dislocation is
 - 1) Inferior dislocation
 - 2) Superior dislocation
 - 3) + Anterior dislocation
 - 4) Posterior dislocation
- 27) The most common type of shock is
 - 1) Septic shock
 - 2) + Hemorrhagic shock
 - 3) Cardiogenic shock
 - 4) Anaphylactic shock
- 28) The medical treatment of hydatid cyst is
 - 1) Metronidazole
 - 2) + Albendazole
 - 3) Ceftriaxone
 - 4) Penecillins
- 29) Two or more contiguous rib fractures with two or more breaks per rib means
 - 1) Pneumthorax
 - 2) + Flial chest
 - 3) Lung cancer

4/ 3 الصفحة



- 4) Chylothorax
- 30) All the followig are part of local examination EXCEPT
 - 1) Inspection
 - 2) Palpation
 - 3) Percussion
 - 4) + Vital signs
- 31) The number of the thoracic vertebrae are
 - 1) 10
 - 2) 11
 - 3) + 12
 - 4) 13
- 32) Battle's sign means there is
 - 1) Anterior cranial fossa fracture
 - 2) + Middle cranial fossa fracture
 - 3) Posterior cranial fossa fracture
 - 4) Fracture of the skull vault
- 33) The most common intra-abdomial solid organ to be injured in blunt trauma is
 - 1) Liver
 - 2) Kidneys
 - 3) + Spleen
 - 4) Stomach
- 34) Erythema of the epidermis is considered
 - 1) + First degree burn
 - 2) Second degree burn
 - 3) Third degree burn
 - 4) Fourthdegree burn
- 35) All the following are part of the lethal triad of hemorrhagic shock EXCEPT
 - 1) Hypothermia
 - 2) Metabolic Acidosis
 - 3) + Metabolic Alkalosis
 - 4) Coagulopathy