



قائمة الاسئلة

علم التشريح التطبيقي - علاج طبيعي - المستوى الثاني - كلية الطب والعلوم الصحية - برامج العلوم الطبية التطبيقية - الفترة الث
د/ صالح عبدالخالق سعد المعرشي

- 1) 1. Pronator teres nerve supply is
 - 1) - a) Radial n
 - 2) + b) Median n
 - 3) - c) Ulnar n
 - 4) - d) None
- 2) 2. All the following are contents of cubital fossa except
 - 1) - a) Radial n
 - 2) - b) Median n
 - 3) + c) Ulnar n
 - 4) - d) Bronchial artery
- 3) 3. Deltoid muscle is supplied by
 - 1) - a) Suprascapular n
 - 2) - b) Pectoral n
 - 3) - c) Radial n
 - 4) + d) Axillary n
- 4) 4. Axillary artery is giving
 - 1) - a) 2 branches
 - 2) - b) 3 branches
 - 3) - c) 4 branches
 - 4) + d) 6 branches
- 5) 5. Total numbers of muscles of the hand are
 - 1) - a) 10 muscles
 - 2) - b) 15 muscles
 - 3) + c) 20 muscles
 - 4) - d) 25 muscles
- 6) 6. All muscles are under the covering of gluteus maximus except
 - 1) - a) Gluteus Medius
 - 2) - b) Gluteus Minimus
 - 3) - c) Piriformis
 - 4) + d) Rectus femoris
- 7) 7. Roots of obturator nerve are
 - 1) - a) L1, L2, L3
 - 2) + b) L2, L3, L4
 - 3) - c) L3, L4, L5
 - 4) - d) L4, L5, S1
- 8) 8. Hamstring muscles are supplied by
 - 1) - a) Femoral nerve
 - 2) - b) Obturator nerve
 - 3) + c) Sciatic nerve
 - 4) - d) None
- 9) 9. Femoral artery gives
 - 1) - a) 3 branches
 - 2) - b) 4 branches
 - 3) - c) 5 branches
 - 4) + d) 6 Branches
- 10) 10. floor of femoral triangle is formed by the following muscles except





- 1) - a) adductor longus
2) - b) psoas major
3) - c) iliacus
4) + d) adductor magnus
- 11) 11. anterior Cruciate ligament is present in
1) - a) hip joint
2) + b) knee joint
3) - c) ankle joint
4) - d) none
- 12) 12. superolateral border of popliteal fossa is formed by
1) + a) biceps femoral m
2) - b) semimembranosus m
3) - c) semitendinosus m
4) - d) gastrocnemius m
- 13) 13. lateral longitudinal arch of the foot is Femoris by
1) + a) 4 bones
2) - b) 5 bones
3) - c) 6 bones
4) - d) 9 bones
- 14) 14. perforator veins are
1) + a) 4 groups
2) - b) 3 groups
3) - c) 5 groups
4) - d) 2 groups
- 15) 15. Dorsal venous arch is formed by
1) - a) 4 Veins
2) + b) 5 veins
3) - c) 3 veins
4) - d) 6 veins
- 16) 16. Calf muscles of the leg is supplied by
1) - a) Common peroneal nerve
2) + b) Tibial nerve
3) - c) Femoral nerve
4) - d) None
- 17) 17. The numbers of tarsal bones
1) - a) 8 bones
2) + b) 7 bones
3) - c) 6 bones
4) - d) 5 bones
- 18) 18. The number of carpal bones are
1) + a) 8 bones
2) - b) 7 bones
3) - c) 6 bones
4) - d) 5 bones
- 19) 19. Small saphenous vein is
1) - a) Deep vein
2) + b) Superficial vein
3) - c) Perforator vein
4) - d) None





- 20) 20. Causes of varicose vein of lower limb is
- 1) + a) Dilatation of superficial vein
 - 2) - b) Dilatation of superficial artery
 - 3) - c) Dilatation of superficial nerve
 - 4) - d) none

