



قائمة الاسئلة

علاج طبيعي لامراض الجراحه و الحروق - ()- المستوى الرابع - قسم - العلاج الطبيعي - كلية الطب والعلوم الصحية - الفترة الأولى- درجة الامتحان

د خالد يحيى السباني

- 1) Skin graft contains its own vasculature
  - 1) - true.
  - 2)  false.
- 2) Epidermis is thicker connective tissue layer
  - 1) - true.
  - 2)  false.
- 3) Hypertrophic scars are a common complication of burn injuries
  - 1)  true.
  - 2) - false.
- 4) Superficial partial thickness burn will require surgical
  - 1) - true.
  - 2)  false.
- 5) In stage III in bedsores of involves the full thickness of the skin
  - 1)  true.
  - 2) - false.
- 6) Cosmetic Surgery: Performed on normal structures of the body to improve appearance
  - 1)  true.
  - 2) - false.
- 7) Psoriasis can be passed from person to person
  - 1) - true.
  - 2)  false.
- 8) With psoriasis, new cells grow every 3 to 4 days
  - 1)  true.
  - 2) - false.
- 9) Pressure Ulcers is a skin disorder that causes skin cells to multiply up to 10 times faster than normal
  - 1) - true.
  - 2)  false.
- 10) In Kleinert Protocol Discontinue splint from 6 week
  - 1)  true.
  - 2) - false.
- 11) Surgery for burned patients is not normally indicated until 48 hours after injury
  - 1)  true.
  - 2) - false.
- 12) A healthy diet is not important in preventing skin breakdown and in aiding wound healing
  - 1)  true.
  - 2) - false.
- 13) Acne is characterized by the formation of comedones
  - 1)  true.
  - 2) - false.
- 14) Physiotherapy is often recommended after urological surgeries to help improve bladder control and reduce pelvic floor dysfunction..
  - 1)  true.
  - 2) - false.
- 15) Physiotherapy has no role in the management of patients following bladder or kidney surgery, as these surgeries do not affect the musculoskeletal system.





- 1) - true.  
2)  + false.
- 16) Physiotherapy is often recommended after urological surgeries to help improve bladder control and reduce pelvic floor dysfunction.  
1)  + true.  
2) - false.
- 17) Early mobilization of the hand is not recommended after extensor tendon repair because it may disrupt the tendon healing process.  
1) - true.  
2)  + false.
- 18) Splinting is commonly used in the early stages of rehabilitation following an extensor tendon repair to protect the tendon and prevent excessive movement.  
1)  + true.  
2) - false.
- 19) After abdominal surgery, physiotherapy is essential for helping prevent deep vein thrombosis (DVT) by encouraging early mobilization and performing leg exercises.  
1)  + true.  
2) - false.
- 20) In the post-operative stage after abdominal surgery, physiotherapy focuses only on breathing exercises, and there is no need for exercises to restore core strength.  
1) - true.  
2)  + false.
- 21) Autograft is  
1) - Donor skin  
2) - Animal skin  
3)  + Own skin  
4) - Cultured skin
- 22) All these are physiotherapy aims for burn patient except :  
1)  + Decrease Strength  
2) - Prevent respiratory complications  
3) - Control Oedema  
4) - Maintain Joint ROM
- 23) Neck in extension is recommended position for :  
1) - Axilla burn  
2) - Posterior neck burn  
3) - Head burn  
4)  + Anterior neck burn
- 24) Which is Cells in the epidermis:  
1) - fibroblasts  
2)  + keratinocytes  
3) - macrophages  
4) - mast cells
- 25) over carpal tunnel is :  
1) - Zone 1  
2) - Zone 2  
3) - Zone 3  
4)  + Zone 4
- 26) The sagittal band splint is used to limit ----- joint flexion of the involved finger :  
1)  + MP





- 2) - DIP  
3) - PIP  
4) - Wrist
- 27) A period of ----- days immobilisation is required to enable graft healing
- 1)  3\5  
2) - 5\7  
3) - 7\9  
4) - 9\10
- 28) In bedsores is the deepest to the epidermis extending into, but no deeper than, the dermis.
- 1) - Stage I  
2)  Stage II  
3) - Stage III  
4) - Stage IV
- 29) All these cardiovascular changes in burn patient except
- 1) - Hypovolemia  
2) - Myocardial depression  
3)  Hypermetabolism  
4) - Edema formation
- 30) Normally, skin cells are replaced every ---- to ---- days.
- 1) - 1 to 10  
2) - 10 to 20  
3)  10 to 30  
4) - 30 to 50
- 31) a chronic progressive inflammatory joint disease that can be associated with psoriasis
- 1) - Spondylitis  
2)  Psoriatic arthritis  
3) - osteoarthritis  
4) - rheumatoid arthritis
- 32) These aims Physiotherapy for bedsores except
- 1) - Reduce edema  
2)  Decrease circulation  
3) - Reduce Pain  
4) - Prevent adherence of wound to underlying tissues
- 33) which occur over the PIP joints in extensor tendon injury
- 1) - Zone 1  
2) - Zone 2  
3)  Zone 3  
4) - Zone 4
- 34) Dynamic splinting in extensor tendon repair it used to the end of
- 1) - 2 weeks  
2) - 4 weeks  
3) - 6 weeks  
4)  8 weeks
- 35) major factors are involved in the pathogenesis of Acne except
- 1)  decreased sebum production  
2) - an abnormality of the microbial flora  
3) - cornification of the pilosebaceous duct  
4) - the production of inflammation
- 36) Acne is most common in





- 1) - back
  - 2)  face
  - 3) - chest
  - 4) - upper limbs
- 37) Which the muscle is effect in Nephrectomy
- 1) - Iliopsoas
  - 2) - Traizups
  - 3)  Latissimus dorsi
  - 4) - extensor of spine
- 38) In urosurgeries Strenuous activity and heavy lifting should be avoided for
- 1) - 2 weeks
  - 2) - 4 weeks
  - 3)  6 weeks
  - 4) - 8 weeks
- 39) Homan's sign is test for
- 1)  DVT
  - 2) - Uremia
  - 3) - hernia
  - 4) - Infection
- 40) Following extensor tendon repair in Zones VIII inpatients are placed in a static volar splint with the wrist in ..... degrees of extension
- 1) - 20
  - 2)  30
  - 3) - 40
  - 4) - 50

