



قائمة الاسئلة 2025-04-09 06:16

الجراحة العامة 2 - () - المستوى الرابع - قسم تخدير - تخدير - كلية الطب والعلوم الصحية - الفترة الأولى - درجة الامتحان (70)

د. وليد المخلافي

1) Which of the following muscles divides the neck into anterior and posterior triangles

- 1) - The anterior digastric muscle
- 2) - The posterior diagastric muscle
- 3) + The sternomastoid muscle
- 4) - The strap muscles

2) The most common cause of acute pancreatitis is

- 1) + Gallbladder stones
- 2) - Idiopathic
- 3) - Drugs
- 4) - Viral infection

3) All the following are clinical features of obstructive jaundice EXCEPT

- 1) + Yellow discoloration of the urine
- 2) - Yellow discoloration of the skin
- 3) - Red discoloration of the urine
- 4) - Pale stool

4) The most common benign lesion of the liver is

- 1) - Hepatic adenoma
- 2) - Hepatic hydatid cyst
- 3) + Hepatic hemangioma
- 4) - Focal nodular hyperplasia

5) All the following are true regarding pyogenic liver abscess EXCEPT

- 1) - It occurs as small lesions in both lobes of the liver
- 2) + It is caused by echinococcus granulosus
- 3) - It is caused by ascending cholangitis
- 4) - It is treated by antibiotics and drainage

6) All the following are triangles of the anterior neck triangle EXCEPT

- 1) - Digastric triangle
- 2) - Carotid triangle
- 3) - Muscular triangle
- 4) + Supraclavicular triangle

7) The spleen is found in the

- 1) + Left hypochondrium
- 2) - Right iliac fossa
- 3) - Left iliac fossa
- 4) - Right hypochondrium

8) All the following are blood supply of the thyroid gland EXCEPT

- 1) - Superior thyroid artery
- 2) + Middle thyroid artery
- 3) - Inferior thyroid artery
- 4) - Thyroidea ima artery

9) The vacuum drain is used after which of the following operations

- 1) + Thyroidectomy
- 2) - Appendectomy
- 3) - Cholecystectomy
- 4) - Splenectomy



- 10) The most common solid abdominal organ to be injured in blunt abdominal trauma is
- 1) - The liver
 - 2) - The small bowel
 - 3) + The spleen
 - 4) - The kidneys
- 11) The most common site of the tip of the appendix is
- 1) - Subcecal
 - 2) - Pre-ileal
 - 3) - Pelvic
 - 4) + Retrocecal
- 12) All the following are true regarding Duodenal ulcer EXCEPT
- 1) - It is more common than gastric ulcer
 - 2) + It carries high risk for malignancy
 - 3) - It occurs in ages younger than gastric ulcer
 - 4) - Biopsy is not necessary to be taken
- 13) When the TSH level is low the patient is in
- 1) + Hyperthyroidism
 - 2) - Euthyroid
 - 3) - Hypothyroidism
 - 4) - Normal thyroid level
- 14) Which of the following sutures is synthetic, monofilament and non absorbable
- 1) - Silk
 - 2) + Prolene
 - 3) - Vicryl
 - 4) - PDS
- 15) All the following are branches of the facial nerve EXCEPT
- 1) - Temporal branch
 - 2) + Maxillary branch
 - 3) - Marginal mandibular branch
 - 4) - Zygomatic branch
- 16) Which of the following is considered as an emergency:
- 1) - Sub-hyoid bursitis
 - 2) + Ludwig Angina
 - 3) - Carotid body tumor
 - 4) - Thyroid cancer
- 17) All the following are predisposing factors for hernia EXCEPT
- 1) - Chronic cough
 - 2) - Straining
 - 3) - Frequent pregnancy
 - 4) + Diarrhea
- 18) The cannula that is size 18G
- 1) - Pink
 - 2) + Green
 - 3) - Grey
 - 4) - Blue
- 19) The thyroid cancer that is familial is the
- 1) - Papillary thyroid cancer
 - 2) + Medullary thyroid cancer
 - 3) - Anaplastic thyroid cancer



- 4) - Follicular thyroid cancer
- 20) The worst serious complication of hernia is:
- 1) - Irreducibility
 - 2) + Strangulation
 - 3) - Incarceration
 - 4) - Obstruction
- 21) All the following investigations are related to the parathyroid gland EXCEPT
- 1) - Calcium level
 - 2) + Postassium level
 - 3) - Phosphate level
 - 4) - Parathyroid hormoe
- 22) The most common swelling in the neck is
- 1) - Thyroid gland enlargment
 - 2) + Lymph node enlargment
 - 3) - Cold abscess
 - 4) - Branchial cyst
- 23) Birds beak appearance on barium swallow means
- 1) - Esophageal carcinoma
 - 2) + Achalasia
 - 3) - Zenker diverticulum
 - 4) - Schatzki's ring
- 24) All the following are true regarding rectus hematoma EXCEPT:
- 1) - This is collection of blood in the rectus sheath
 - 2) - It mostly occurs after trauma or may occur spontaneously in patient on anticoagulants
 - 3) + There is a male predominance
 - 4) - The bleeding comes mostly from the inferior epigastric artery
- 25) Surgical removal of the gallbladder is called
- 1) - Coloectomy
 - 2) + Cholecystectomy
 - 3) - Choledolithotomy
 - 4) - Splenectomy
- 26) Which of the following umbilical lesions is due to hair and can appear in other sites like the lower back:
- 1) - Umbilical fistula
 - 2) - Umbilical abscess
 - 3) + Umbilical pilonidal sinus
 - 4) - Umbilical hernia
- 27) The following neck swellings occur in the carotid triangle EXCEPT
- 1) - Carotid body tumor
 - 2) - Branchial cyst
 - 3) + Thyroglossal cyst
 - 4) - Lymph node
- 28) The operation for hernia where the surgeon uses mesh is called
- 1) - Herniotomy
 - 2) - Herniorraphy
 - 3) + Hernioplasty
 - 4) - Herniectomy
- 29) All the following are requirements for incisions EXCEPT:
- 1) + Compressibility
 - 2) - Accessibility





- 3) - Safety
4) - Extensibility
- 30) Which of the following swelling moves with protrusion (movement) of the tongue
- 1) - Dermoid cyst
2) - Lymph node
3) + Thyroglossal cyst
4) - Lipoma
- 31) The instrument used to hold the needle during suturing is
- 1) - Artery forceps
2) + Needle holder
3) - Non-tooth forceps
4) - Toothed forceps
- 32) The suture technique that is cosmetic (good looking) is called
- 1) + Subcuticular suturing
2) - Interrupted suturing
3) - Vertical mattress suturing
4) - Horizontal mattress suturing
- 33) The sigmoid is the most common site for the following diseases EXCEPT
- 1) - Colon cancer
2) - Volvulus
3) - Diverticulosis
4) + Ischemic colitis
- 34) Which of the following muscle is not part of the anterior abdominal wall
- 1) - External oblique muscle
2) - Transversus abdominus muscle
3) + Pectoralis major muscle
4) - Internal oblique muscle
- 35) All the following are true regarding carotid body tumor EXCEPT
- 1) - It occurs in the carotid triangle of the neck
2) - Mostly treated by surgery
3) - It arises at the bifurcation of the common carotid artery
4) + It arises from the baroreceptors