



قائمة الاسئلة 2025-04-13 09:02

علم الادوية - (0) - المستوى الثاني - قسم جميع التخصصات - كلية العلوم الطبية التطبيقية - الفترة الثانية - درجة الامتحان (70)

أ.د. عدنان الادهل

- 1) The drug used for myasthenia gravis is:
 - 1) - adrenaline
 - 2) - dobutamine
 - 3) - salbutamol
 - 4) ☒ neostigmine
 - 5) - propranolol
- 2) The drug used for asthma is:
 - 1) - adrenaline
 - 2) - dobutamine
 - 3) ☒ salbutamol
 - 4) - neostigmine
 - 5) - propranolol
- 3) The drug used for acute heart failure is:
 - 1) - adrenaline
 - 2) ☒ dobutamine
 - 3) - salbutamol
 - 4) - neostigmine
 - 5) - propranolol
- 4) The drug used for anaphylactic shock is:
 - 1) ☒ adrenaline
 - 2) - dobutamine
 - 3) - salbutamol
 - 4) - neostigmine
 - 5) - propranolol
- 5) The drug used for migraine prophylaxis is:
 - 1) - adrenaline
 - 2) - dobutamine
 - 3) - salbutamol
 - 4) - neostigmine
 - 5) ☒ propranolol
- 6) The best drug among the following for morning sickness is:
 - 1) - chlorpheniramine
 - 2) - diphenhydramine
 - 3) ☒ meclizine plus vitamin B6
 - 4) - cyproheptadine
 - 5) - cetirizine
- 7) The best drug among the following for motion sickness is:
 - 1) - chlorpheniramine
 - 2) ☒ diphenhydramine
 - 3) - meclizine plus vitamin B6
 - 4) - cyproheptadine
 - 5) - cetirizine
- 8) The best drug among the following for anaphylaxis is:
 - 1) ☒ chlorpheniramine
 - 2) - diphenhydramine



- 3) - meclizine plus vitamin B6
- 4) - cyproheptadine
- 5) - cetirizine
- 9) The best drug among the following for migraine prophylaxis is:
- 1) - chlorpheniramine
- 2) - diphenhydramine
- 3) - meclizine plus vitamin B6
- 4) + cyproheptadine
- 5) - cetirizine
- 10) The non-sedating drug is:
- 1) - chlorpheniramine
- 2) - diphenhydramine
- 3) - meclizine plus vitamin B6
- 4) - cyproheptadine
- 5) + cetirizine
- 11) The best drug among the following for anaphylaxis is:
- 1) + hydrocortisone
- 2) - prednisolone
- 3) - methylprednisolone
- 4) - beclomethasone
- 5) - betamethasone
- 12) The best drug among the following for acute autoimmune diseases is:
- 1) - hydrocortisone
- 2) - prednisolone
- 3) + methylprednisolone
- 4) - beclomethasone
- 5) - betamethasone
- 13) The best drug among the following for chronic autoimmune diseases is:
- 1) - hydrocortisone
- 2) + prednisolone
- 3) - methylprednisolone
- 4) - beclomethasone
- 5) - betamethasone
- 14) The best drug among the following for topical application is:
- 1) - hydrocortisone
- 2) - prednisolone
- 3) - methylprednisolone
- 4) - beclomethasone
- 5) + betamethasone
- 15) The best drug among the following for asthma is:
- 1) - hydrocortisone
- 2) - prednisolone
- 3) - methylprednisolone
- 4) + beclomethasone
- 5) - betamethasone
- 16) The best anti-inflammatory drug among the following for children is:
- 1) - aspirin
- 2) - paracetamol
- 3) + ibuprofen



- 4) - diclofenac
5) - mefenamic acid
- 17) The most powerful anti-inflammatory drug among the following is:
1) - aspirin
2) - paracetamol
3) - ibuprofen
4) ☒ diclofenac
5) - mefenamic acid
- 18) The best anti-inflammatory drug among the following for dysmenorrhoea is:
1) - aspirin
2) - paracetamol
3) - ibuprofen
4) - diclofenac
5) ☒ mefenamic acid
- 19) The antiplatelet drug is:
1) ☒ aspirin
2) - paracetamol
3) - ibuprofen
4) - diclofenac
5) - mefenamic acid
- 20) The anti-inflammatory drug that is safe for patients with peptic ulcer is:
1) - aspirin
2) ☒ paracetamol
3) - ibuprofen
4) - diclofenac
5) - mefenamic acid
- 21) The opioid that is used for severe pain is:
1) ☒ morphine
2) - codeine
3) - loperamide
4) - naloxone
5) - naltrexone
- 22) The opioid that is used for moderate pain is:
1) - morphine
2) ☒ codeine
3) - loperamide
4) - naloxone
5) - naltrexone
- 23) The opioid that is used for non-infectious diarrhoea is:
1) - morphine
2) - codeine
3) ☒ loperamide
4) - naloxone
5) - naltrexone
- 24) The opioid antagonist that is used for opioids overdose is:
1) - morphine
2) - codeine
3) - loperamide
4) ☒ naloxone





- 5) - naltrexone
- 25) The opioid antagonist that is used for the treatment of opioids addicts is:
- 1) - morphine
- 2) - codeine
- 3) - loperamide
- 4) - naloxone
- 5) + naltrexone
- 26) The best anti-spasticity among the following is:
- 1) - diazepam
- 2) + baclofen
- 3) - tizanidine
- 4) - dantrolene
- 5) - botulinum toxin
- 27) Hypotension is a side effect most likely occur with the use of:
- 1) - diazepam
- 2) - baclofen
- 3) + tizanidine
- 4) - dantrolene
- 5) - botulinum toxin
- 28) Sedation is a side effect most likely occur with the use of:
- 1) + diazepam
- 2) - baclofen
- 3) - tizanidine
- 4) - dantrolene
- 5) - botulinum toxin
- 29) Skeletal muscle weakness is a side effect most likely occur with the use of:
- 1) - diazepam
- 2) - baclofen
- 3) - tizanidine
- 4) - dantrolene
- 5) + botulinum toxin
- 30) The drug that inhibit the release of calcium from sarcoplasmic reticulum is:
- 1) - diazepam
- 2) - baclofen
- 3) - tizanidine
- 4) + dantrolene
- 5) - botulinum toxin
- 31) Polyuria is a side effect most likely occur with the use of:
- 1) - glibenclamide
- 2) - metformin
- 3) + empagliflozin
- 4) - acarbose
- 5) - candesartan
- 32) Weight loss is a side effect most likely occur with the use of:
- 1) - glibenclamide
- 2) + metformin
- 3) - empagliflozin
- 4) - acarbose
- 5) - candesartan



- 33) Weight gain is a side effect most likely occur with the use of:
- 1) ☒ glibenclamide
 - 2) ☐ metformin
 - 3) ☐ empagliflozin
 - 4) ☐ acarbose
 - 5) ☐ candesartan
- 34) Diarrhoea is a side effect most likely occur with the use of:
- 1) ☐ glibenclamide
 - 2) ☐ metformin
 - 3) ☐ empagliflozin
 - 4) ☒ acarbose
 - 5) ☐ candesartan
- 35) Hyperkalaemia is a side effect most likely occur with the use of:
- 1) ☐ glibenclamide
 - 2) ☐ metformin
 - 3) ☐ empagliflozin
 - 4) ☐ acarbose
 - 5) ☒ candesartan
- 36) The antiviral drug that is used for herpes simplex virus is:
- 1) ☐ Benzathine penicillin
 - 2) ☐ fluconazole
 - 3) ☒ acyclovir
 - 4) ☐ albendazole
 - 5) ☐ metronidazole
- 37) The antifungal drug that is used for tinea infection is:
- 1) ☐ Benzathine penicillin
 - 2) ☒ fluconazole
 - 3) ☐ aciclovir
 - 4) ☐ albendazole
 - 5) ☐ metronidazole
- 38) The anthelmintic drug that is used for ascariasis is:
- 1) ☐ Benzathine penicillin
 - 2) ☐ fluconazole
 - 3) ☐ aciclovir
 - 4) ☒ albendazole
 - 5) ☐ metronidazole
- 39) The antiprotozoal drug that is used for amoebiasis is:
- 1) ☐ Benzathine penicillin
 - 2) ☐ fluconazole
 - 3) ☐ aciclovir
 - 4) ☐ albendazole
 - 5) ☒ metronidazole
- 40) The antibacterial drug that is used for rheumatic heart fever is:
- 1) ☒ Benzathine penicillin
 - 2) ☐ fluconazole
 - 3) ☐ aciclovir
 - 4) ☐ albendazole
 - 5) ☐ metronidazole