

قائمة الاسئلة 04:16 00-2025

العلاج الطبيعي لأمراض الجهاز العضلي والهيكلي2-الرابع- علاج طبيعي-كلية الطب-درجةالامتحان (80)

د/ خالد يحيى السياني

- 1) A muscle strength grade of 2 indicates active movement with gravity eliminated.
 - 1) + .TRUE
 - 2) .FALSE
- 2) The Adam's forward bend test is used to diagnose thoracic outlet syndrome.
 - 1) .TRUE
 - 2) + .FALSE
- 3) The Spurling test is used to diagnose cervical spondylosis.
 - 1) .TRUE
 - 2) + .FALSE
- 4) torticollis referred to as cervical dystonia
 - 1) + .TRUE
 - 2) .FALSE
- 5) Cervical flexion occurs in the sagittal plane.
 - 1) + .TRUE
 - 2) .FALSE
- 6) Spondylolisthesis is most often caused by spondylolysis
 - 1) + .TRUE
 - 2) .FALSE
- 7) An important part of the diagnosis of low back pain includes palpation of the lumbar spinous processes.
 - 1) + .TRUE
 - 2) .FALSE
- 8) horacic outlet syndrome is associated with compression of the brachial plexus and subclavian artery.
 - 1) + .TRUE
 - 2) .FALSE
- 9) The lumbar spine allows only flexion and extension movements.
 - 1) .TRUE
 - 2) + .FALSE
- 10) Osteoarthritis is the most common form of arthritis, and the hip is the frist most commonly affected joint
 - 1) .TRUE
 - 2) + .FALSE
- 11) The intervertebral disc consists of an outer annulus fibrosus and an inner nucleus pulposus.
 - 1) + .TRUE
 - 2) .FALSE
- 12) Physical therapy cannot treatment avascular necrosis
 - 1) + .TRUE
 - 2) .FALSE
- Degenerative Disc Disease (DDD) is primarily caused by bacterial infections in the spine.
 - 1) .TRUE
 - 2) + .FALSE
- 14) The knee joint is the biggest joint in the body
 - 1) + .TRUE
 - 2) .FALSE
- 15) Spondylosis is a type of arthritis that affects the spine.
 - 1) + .TRUE
 - 2) .FALSE

4 / 1 الصفحة



- Pre-operative of TKR the patient not need physical therapy session
 - 1) .TRUE
 - 2) + .FALSE
- 17) Knee osteoarthritis (OA) is the most common cause of disability in adults.
 - 1) + .TRUE
 - 2) .FALSE
- 18) In Day of Surgery TKR use CPM 0-140° started in Recovery Room for minimum of 4 hours
 - 1) 1
 - 2) + 0
- 19) The Lachman test is used to assess meniscus injuries.
 - 1) .TRUE
 - 2) + .FALSE
- 20) Muscle strains most commonly occur in bi-articular muscles such as the hamstrings, rectus femoris and gastrocnemius
 - 1) + 1
 - 2) 0
- 21) Which of the following is NOT a typical symptom of cervical radiculopathy?
 - 1) Neck pain
 - 2) Radiating arm pain
 - 3) + Shortness of breath
 - 4) Muscle weakness
- 22) What is the first-line treatment for cervical disc prolapse?
 - 1) Surgery
 - 2) + NSAIDs and physical therapy
 - 3) Botulinum toxin injection
 - 4) Complete immobilization
- 23) What is the correct patient position when testing cervical flexion with a goniometer?
 - 1) + Sitting with thoracic and lumbar spine supported
 - 2) Standing with arms extended
 - 3) Lying supine with neck relaxed
 - 4) Prone with arms behind the back
- 24) What is the natural curvature of the thoracic spine?
 - 1) Lordosis
 - 2) Scoliosis
 - 3) Neutral
 - 4) + Kyphosis
- 25) Milwaukee brace is use for
 - 1) + Scheuermann's disease
 - 2) cervical rib
 - 3) disc prolapse
 - 4) spondylosis
- 26) All these Symptoms of Degenerative Disc Disease of lumber spine except
 - 1) Tingling or numbness in the lower limb
 - 2) Stiffness
 - 3) + Weakness in the upper limb
 - 4) Pain in the lower back
- 27) What is the most common cause of kyphotic deformity in adolescents?
 - 1) Cervical rib
 - 2) Osteoporosis

4 / 2 الصفحة



- 3) Scoliosis
- 4) + Scheuermann's disease
- 28) Which nerve root is tested by ankle dorsiflexion strength?
 - 1) L2
 - 2) L3
 - + L4
 - 4) L5
- 29) Which treatment is NOT commonly used for degenerative disc disease?
 - 1) Nonsteroidal anti-inflammatory drugs (NSAIDs)
 - 2) + Chemotherapy
 - 3) Physical therapy
 - 4) Steroid injections
- 30) Passive lumbar extension test use for :
 - 1) Lumber Facet joint pathology
 - 2) + Lumbar instability
 - 3) Spondylolisthesis
 - 4) neurological dysfunction
- 31) Which is Not risk factors for OA
 - 1) + Young age
 - 2) Family history of osteoarthritis
 - 3) Previous injury to the hip joint
 - 4) Obesity
- 32) All these Goals management of OA except
 - 1) Patient education
 - 2) Muscle strengthening
 - 3) + Decrease ROM
 - 4) Modification of risk factors
- 33) During hip joint palpation, which structure is commonly assessed for tenderness?
 - 1) + Greater trochanter
 - 2) Medial malleolus
 - 3) Clavicle
 - 4) Tibial tuberosity
- What is the most sensitive imaging modality for detecting early avascular necrosis?
 - 1) X-ray
 - 2) Ultrasound
 - 3) + MRI
 - 4) CT scan
- Which special test is used to assess posterior cruciate ligament (PCL) injury?
 - 1) Lachman test
 - 2) Anterior drawer test
 - 3) + Posterior drawer test
 - 4) McMurray test
- 36) Genu varum refers to:
 - 1) + Bowed legs (knees bending outward)
 - 2) Knock-knees (knees bending inward)
 - 3) Hyperextension of the knee
 - 4) Locked knee joint
- 37) All these goals Physical therapy tretment for Avascular Necrosis of the Hip except
 - 1) Relief of symptoms



- 2) + Increase of muscles weakness
- 3) Prevention of disease progression
- 4) Improvement of functionality
- 38) What is a common complication after total knee arthroplasty (TKA)?
 - 1) Avascular necrosis of the hip
 - 2) + Deep vein thrombosis (DVT)
 - 3) Cervical radiculopathy
 - 4) Carpal tunnel syndrome
- 39) A positive Thompson test indicates:
 - 1) + Achilles tendon rupture
 - 2) Plantar fasciitis
 - 3) Tibial nerve compression
 - 4) Lateral ligament sprain
- 40) Which is NOT Physiotherapy Goals post Total Knee Replacement operation
 - 1) Optimise respiratory and circulatory functions
 - 2) Facilitate recovery of quads strength
 - 3) + Increase post op oedema
 - 4) Promote early ambulation